bsm-52166R

[Primary Antibody]

phospho-HSF1 (Ser326) Recombinant Rabbit **mAb**



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG Clonality: Recombinant CloneNo.: 37E4 **GenelD: 3297 SWISS:** Q00613

Target: HSF1 (Ser326)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

HSF1 around the phosphorylation site of Ser326: L(p-S)PT.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: Size: 25ul/50ul/100ul/200ul

0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Size: 200ug (PBS only)

0.01M PBS

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The product of this gene is a heat-shock transcription factor.

Transcription of heat-shock genes is rapidly induced after temperature stress. Hsp90, by itself and/or associated with multichaperone complexes, is a major repressor of this gene.

[provided by RefSeq, Jul 2008].

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (2ug/Test)

Reactivity: Human, Mouse


(predicted: Rat)

Predicted MW.:


Subcellular

Location: Cytoplasm ,Nucleus


VALIDATION IMAGES




Sample: Lane 1: Normal human HeLa cell lysates Lane 2: Hela cells heated at 42 °C for 30 minutes Primary: Anti-phospho-HSF1 (Ser326) (bsm-52166R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 57 kDa Observed band size: 80 kDa



Paraformaldehyde-fixed, paraffin embedded (human endometrial carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-HSF1 (Ser326)) Polyclonal Antibody, Unconjugated (bsm-52166R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min: Antibody incubation with (phospho-HSF1 (Ser326)) Monoclonal Antibody, Unconjugated (bsm-52166R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control:MCF7. Primary Antibody (green line): Rabbit Anti-phospho-HSF1 (Ser326) antibody (bsm-52166R) Dilution: 2µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF488 Dilution: $1\mu g$ /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at-20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.