

bsm-52171R**[Primary Antibody]**

BioSS
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

phospho-JAK2 (Tyr1007+Tyr1008) Recombinant Rabbit mAb


— DATASHEET —

Host: Rabbit Clonality: Recombinant GeneID: 3717 Target: phospho-JAK2 (Tyr1007+Tyr1008) Immunogen: A synthesized peptide derived from human JAK2 around the phosphorylation site of Y1007/Y1008: KE-pY-pY-KV. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: This gene product is a protein tyrosine kinase involved in a specific subset of cytokine receptor signaling pathways. It has been found to be constitutively associated with the prolactin receptor and is required for responses to gamma interferon. Mice that do not express an active protein for this gene exhibit embryonic lethality associated with the absence of definitive erythropoiesis. [provided by RefSeq, Jul 2008]	Isotype: IgG CloneNo.: 6E5 SWISS: O60674	Applications: IHC-P (1:50-200) IHC-F (1:50-200) IF (1:50-200) ICC/IF (1:50-200) Reactivity: Human, Mouse, Rat Predicted MW.: 131 kDa Subcellular Location: Cell membrane ,Cytoplasm ,Nucleus
--	---	--


— VALIDATION IMAGES —



Paraformaldehyde-fixed, paraffin embedded Mouse Lung; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with phospho-JAK2 (Tyr1007+Tyr1008) Monoclonal Antibody, Unconjugated(bsm-52171R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit(Rabbit, SP-0023) and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Gastric Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with phospho-JAK2 (Tyr1007+Tyr1008) Monoclonal Antibody, Unconjugated(bsm-52171R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit(Rabbit, SP-0023) and DAB (C-0010) staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Phospho-JAK2(Y1007+Y1008)) monoclonal Antibody, Unconjugated (bsm-52171R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

— SELECTED CITATIONS —

- **[IF=8.4]** Hu Bowen. et al. Local GHR roles in regulation of mitochondrial function through mitochondrial biogenesis during myoblast differentiation. CELL COMMUN SIGNAL. 2023 Dec;21(1):1-18 WB ;Chicken. 37337300
- **[IF=2.985]** Yanyan Lu. et al. Silencing IFN γ inhibits A1 astrocytes and attenuates neurogenesis decline and cognitive impairment in endotoxemia. Biochem Bioph Res Co. 2020 Dec;533:1519 WB,IHC ;Mouse. 33158480