bs-17861R

- DATASHEET -

[Primary Antibody]

MSRB2 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Host: Rabbit Isotype: IgG Clonality: Polyclonal IF (1:100-500) GenelD: 22921 SWISS: Q9Y3D2 Target: MSRB2 Immunogen: KLH conjugated synthetic peptide derived from human MSRB2: Horse) 31-130/182. Purification: affinity purified by Protein A Predicted 20 kDa MW.: Concentration: 1mg/ml Subcellular Location: Cytoplasm Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Methionine is one of the most readily oxidized essential amino acids and an intermediate in the biosynthesis of cysteine, carnitine, taurine, lecithin, phosphatidylcholine and other phospholipids. In its oxidative state, Methionine is regulated in vivo by methionine sulfoxide reductases (Msr). MsrB2 is a 182 amino acid mitochondrial protein that is ubiquitously expressed. Belonging to the MsrB Met sulfoxide reductase family, MsrB2 acts as a catalyst for the reduction of free and protein-bound methionine sulfoxide to methionine. Upon oxidative stress, MsrB2 is suggested to play a role in the preservation of mitochondrial integrity by decreasing the intracellular reactive oxygen species build-up through its scavenging role, hence contributing to cell survival and protein maintenance. MsrB2 utilizes zinc ions, one per subunit, as cofactors.

VALIDATION IMAGES



Tissue/cell: mouse embryo tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum.C-0005) at 37°C for 20 min: Incubation: Anti-MSRB2 Polyclonal Antibody, Unconjugated(bs-17861R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum.C-0005) at 37°C for 20 min: Incubation: Anti-MSRB2 Polyclonal Antibody, Unconjugated(bs-17861R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

– SELECTED CITATIONS —

• [IF=2.639] Gaolong Zhong, et al. Effects of Long-Term Exposure to Copper on the Keap1/Nrf2 Signaling Pathway and Msr-Related Redox Status in the Kidneys of Rats. 2021 Jan 22 IF ;Rat. 33479888

Applications: IHC-P (1:100-500) IHC-F (1:100-500)

Reactivity: Mouse, Rat (predicted: Human, Dog,