

bs-3369R**[Primary Antibody]**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Phospho-CDC42 (Ser71) Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: ICC/IF (1:100)
Clonality: Polyclonal		
GeneID: 998	SWISS: P60953	
Target: Phospho-CDC42 (Ser71)		
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human CDC42 around the phosphorylation site of Ser71: PL(p-S)YP.		
Purification: affinity purified by Protein A		Reactivity: Human (predicted: Mouse, Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse)
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 21 kDa
Background: The protein encoded by this gene is a small GTPase of the Rho-subfamily, which regulates signaling pathways that control diverse cellular functions including cell morphology, migration, endocytosis and cell cycle progression. This protein is highly similar to <i>Saccharomyces cerevisiae</i> Cdc 42, and is able to complement the yeast <i>cdc42-1</i> mutant. The product of oncogene <i>Dbl</i> was reported to specifically catalyze the dissociation of GDP from this protein. This protein could regulate actin polymerization through its direct binding to Neural Wiskott-Aldrich syndrome protein (N-WASP), which subsequently activates Arp2/3 complex. Alternative splicing of this gene results in multiple transcript variants. Pseudogenes of this gene have been identified on chromosomes 3, 4, 5, 7, 8 and 20. [provided by RefSeq, Apr 2013]		Subcellular Location: Cytoplasm

— SELECTED CITATIONS —

- **[IF=2.049]** Lei Q et al. HEV ORF3 down - regulates CD14 and CD64 to impair macrophages phagocytosis through inhibiting JAK/STAT pathway. Journal of Medical Virology. 2019. WB ;Human. doi:10.1002/jmv.25400