bs-3380R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

IHC-F (1:100-500)

IF (1:100-500)

Reactivity: Mouse (predicted: Human,

102 kDa

Rat, Rabbit, Chicken)

Applications: IHC-P (1:100-500)

Predicted

MW.:

Subcellular Nucleus

phospho-Rb (Ser807 + Ser811) Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 5925 **SWISS:** P06400

Target: Rb (Ser807 + Ser811)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

Rb around the phosphorylation site of Ser807/Ser811: YI(p-

S)PLK(p-S)PY.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is a negative regulator of the cell cycle and was the first tumor suppressor gene found. The encoded


protein also stabilizes constitutive heterochromatin to maintain the overall chromatin structure. The active, hypophosphorylated form of the protein binds transcription factor E2F1. Defects in this gene are a cause of childhood cancer retinoblastoma (RB), bladder

cancer, and osteogenic sarcoma.

Rb is a tumor suppressor gene which functions as a negative regulator of the cell cycle by interacting with transcription factors including E2F1, PU1, ATF2, UBF, Elf1 and cAbl. This ability of Rb to alter transcription is regulated by phosphorylation catalyzed by the cyclin dependent protein kinases (cdks). Rb is phosphorylated on serine and threonine, but not on tyrosine residues. It forms a complex with SV40 large T antigen, adenovirus E1A, and human papilloma virus 16E. Rb protein may act by regulating transcription and loss of its function leads to uncontrolled cell growth. Aberrations in the Rb gene have been implicated in cancers of

breast, colon, prostate, kidney, nasopharynx, and leukemia.

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse eye); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-Rb (Ser807 + Ser811)) Polyclonal Antibody, Unconjugated (bs-3380R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

- SELECTED CITATIONS -

- [IF=4.183] Zou P et al. Mechanisms of Stress-Induced Spermatogenesis Impairment in Male Rats Following Unpredictable Chronic Mild Stress (uCMS). Int. J. Mol. Sci. 2019, 20, 4470. WB; Rat. doi:10.3390/ijms20184470
- [IF=2.58] Huang, Xu, et al. "ICAM-1-Targeted Liposomes Loaded with Liver X Receptor Agonists Suppress PDGF-Induced Proliferation of Vascular Smooth Muscle Cells." Nanoscale Research Letters 12.1 (2017): 322. WB; Mouse. 28472871
- [IF=3.263] Fei Min et al. Hepatoprotective effects of hydroxysafflor yellow A in D-galactose-treated aging mice. Eur J Pharmacol. 2020 Aug 15;881:173214. WB,IHC; Mouse. 32454116