bs-7496R

[Primary Antibody]

WDR35 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500) IF (1:100-500)
GenelD: 57539	SWISS: Q9P2L0	ICC/IF (1:100-500)
Target: WDR35		ELISA (1:5000-10000)
Immunogen: KLH conjugated synthetic peptide derived from human WDR35: 421-520/1181.		Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Sheep, Cow, Dog)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Prodicted
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: ^{133 kDa} Subcellular Location: ^{Cytoplasm}
repeats are minim amino acids typica which may facilita complexes. Memb cellular processes, transduction, apo alternatively splice have been found fo syndrome / cranio mutations in this g	a member of the WD repeat protein family. WD ally conserved regions of approximately 40 illy bracketed by gly-his and trp-asp (GH-WD), te formation of heterotrimeric or multiprotein ers of this family are involved in a variety of including cell cycle progression, signal otosis, and gene regulation. Multiple ed transcript variants encoding distinct isoforms or this gene. Two patients with Sensenbrenner ectodermal dysplasia (CED) were identified with gene, consistent with a possible ciliary I by RefSeq, Sep 2010]	