## bs-7421R

## [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# Sprouty 4 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GeneID:** 81848 **SWISS:** Q9C004

Target: Sprouty 4

**Immunogen:** KLH conjugated synthetic peptide derived from human Sprouty 4:

1-100/299.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: SPRY4 is an inhibitor of the receptor-transduced mitogen-activated

protein kinase (MAPK) signaling pathway. It is positioned upstream of RAS (see HRAS; MIM 190020) activation and impairs the formation of active GTP-RAS (Leeksma et al., 2002 [PubMed

12027893]).[supplied by OMIM, Mar 2008]

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) ICC/IF (1:100-500) **ELISA** (1:5000-10000)

Reactivity: (predicted: Human, Mouse,

Rat, Cow, Dog, Horse)

Predicted MW.: 33 kDa

Subcellular Location: Cytoplasm

### - SELECTED CITATIONS -

• [IF=31.745] Shan-shan Liu. et al. The chemokine CCL1 triggers an AMFR-SPRY1 pathway that promotes differentiation of lung fibroblasts into myofibroblasts and drives pulmonary fibrosis. Immunity. 2021 Aug;: WB; Mouse. 34407391