

bs-20421R**[Primary Antibody]****COUP-TF1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 7025	SWISS: P10589	
Target: COUP-TF1		Reactivity: Rat (predicted: Human, Mouse, Pig, Sheep, Cow)
Immunogen: KLH conjugated synthetic peptide derived from human COUP-TF1: 21-120/423.		
Purification: affinity purified by Protein A		Predicted MW.: 46 kDa
Concentration: 1mg/ml		Subcellular Location: Nucleus
Storage: Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: COUP transcription factors have been cloned in several species and identified as orphan members of the steroid/thyroid hormone receptor superfamily. COUP-TFI (also designated COUP or EAR-3) and ARP-1 (also designated COUP-TFII) exhibit highly regulated and overlapping expression in most tissues. COUP-TFs are highly expressed in the developing and central nervous system, suggesting that these factors may be important in neural development and differentiation. COUP-TFs can compete for binding to response elements which are common to other members of this family, including RAR, RXR, PPAR, HNF-4, VDR and TR. They have been shown to act as negative regulators as well as initiators of transcription.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat lung tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (COUP-TF1) Polyclonal Antibody, Unconjugated (bs-20421R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.