

---

## ASCL1 Rabbit pAb

Catalog Number: bs-1155R

Target Protein: ASCL1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), Flow-Cyt (2ug/Test), ICC/IF (1:100)

Reactivity: Human, Mouse, Rat (predicted:Sheep, Cow)

Predicted MW: 26 kDa

Entrez Gene: 429

Swiss Prot: P50553

Source: KLH conjugated synthetic peptide derived from human ASCL1: 151-236/236.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** This gene encodes a member of the basic helix-loop-helix (BHLH) family of transcription factors. The protein activates transcription by binding to the E box (5'-CANNTG-3'). Dimerization with other BHLH proteins is required for efficient DNA binding. This protein plays a role in the neuronal commitment and differentiation and in the generation of olfactory and autonomic neurons. Mutations in this gene may contribute to the congenital central hypoventilation syndrome (CCHS) phenotype in rare cases. [provided by RefSeq, Jul 2008]

### VALIDATION IMAGES


---




Sample: Brain (Mouse) Lysate at 40 ug Primary: Anti-ASCL1 (bs-1155R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 26 kD Observed band size: 26 kD




Tissue/cell: human glioma tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-MASH1 Polyclonal Antibody, Unconjugated(bs-1155R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-MASH1 Polyclonal Antibody, Unconjugated(bs-1155R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: mouse embryos tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-MASH1 Polyclonal Antibody, Unconjugated(bs-1155R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-MASH1 Polyclonal Antibody, Unconjugated(bs-1155R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control:Mouse brain. Primary Antibody (green line): Rabbit Anti-MASH1 antibody (bs-1155R) Dilution: 2µg /10<sup>6</sup> cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

## PRODUCT SPECIFIC PUBLICATIONS

[IF=5.41] Misra et al. Asymmetric activation of Dll4-Notch signaling by Foxn4 and proneural factors activates BMP/TGFβ signaling to specify V2b interneurons in the spinal cord. (2014) Development. 141:187-98 ChIP ; Mouse . 24257627

[IF=2.535] Song X et al. Anti-aging effects exerted by Tetramethylpyrazine enhances self-renewal and neuronal differentiation of rat

bMSCs by suppressing NF- $\kappa$ B signaling. Biosci Rep. 2019 Jun 25;39(6). pii: BSR20190761. WB ; Rat . 31171713

[IF=2.886] Xiao Fu. et al. PD-L1 Predicts Poor Prognosis in Surgically Resected Limited Stage Small-Cell Lung Cancer. Cancer Manag Res. 2020; 12: 10939–10948 IHC ; Human . 33154673

[IF=2.6] Li Jianan. et al. Characteristics of molecular subtypes and clinical outcomes in the immunotherapy Queue of extensive-stage small cell lung cancer patients. BMC PULM MED. 2024 Dec;24(1):1-12 IHC ; Human . 39210380

[IF=2.3] Prieto T.G.. et al. Clinical and morphological features of large-cell neuroendocrine carcinomas and small-cell lung carcinomas expressing the DLL3 and ASCL1 oncoproteins. BRAZ J MED BIOL RES. 2023 Dec;56:e12921 IHC ; Human . 38126617