

bs-0682R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

PDK4 Rabbit pAb**— DATASHEET —**


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse, Rat Predicted MW.: 47 kDa Subcellular Location: Cell membrane ,Cytoplasm
Clonality: Polyclonal		
GeneID: 5166	SWISS: Q16654	
Target: PDK4		
Immunogen: KLH conjugated synthetic peptide derived from human Pdk4: 311-411/411.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene is a member of the PDK/BCKDK protein kinase family and encodes a mitochondrial protein with a histidine kinase domain. This protein is located in the matrix of the mitochondria and inhibits the pyruvate dehydrogenase complex by phosphorylating one of its subunits, thereby contributing to the regulation of glucose metabolism. Expression of this gene is regulated by glucocorticoids, retinoic acid and insulin. [provided by RefSeq, Jul 2008].		

— VALIDATION IMAGES —

Sample: Lane 1: Mouse Cerebrum tissue lysates
 Lane 2: Rat Cerebrum tissue lysates
 Primary: Anti-PDK4 (bs-0682R) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 47 kDa
 Observed band size: 45 kDa



Paraformaldehyde-fixed, paraffin embedded (human myocardium); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PDK4) Polyclonal Antibody, Unconjugated (bs-0682R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PDK4) Polyclonal Antibody, Unconjugated (bs-0682R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=6.5]** Yang Fan. et al. Identifying oxidative stress-related biomarkers in idiopathic pulmonary fibrosis in the context of predictive, preventive, and personalized medicine using integrative omics approaches and machine-learning strategies. EPMA Journal. 2023 Jul;:1-26 WB ;Rat. 10.1007/s13167-023-00334-4
- **[IF=6.9]** Liang Song. et al. High-fat intake during lactation ameliorates cardiac fatty acid metabolic disorders and dysfunction in mouse offspring undergoing prenatal poly (I:C) stimulation. ACTA PHARMACOLOGICA SINICA. 2025 Feb 25. Western Blot ;Mouse. 10.1038/s41401-025-01497-8
- **[IF=0]** Zongwang Zhang. et al. AARS2 ameliorates myocardial ischemia via fine-tuning PKM2-mediated metabolism.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

