bsm-60920R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

CD33 Recombinant Rabbit mAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Recombinant CloneNo.: 14H11

GeneID: 945 SWISS: P20138

Target: CD33

Immunogen: A synthesized peptide derived from human CD33: 1-70/364.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS(pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20 °C for one year. Avoid repeated

freeze/thaw cycles.

Background: Enables protein phosphatase binding activity and sialic acid

binding activity. Involved in several processes, including negative regulation of cytokine production; negative regulation of monocyte activation; and positive regulation of protein tyrosine phosphatase activity. Located in several cellular components, including Golgi apparatus; external side of plasma membrane; and peroxisome. [provided by Alliance of Genome Resources, Apr 2022]

Applications: WB (1:500-2000)


IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)

Reactivity: Human


Predicted MW.: 37 kDa

Subcellular Location: Cell membrane


VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with CD33 monoclonal antibody, unconjugated (bsm-60920R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD33 Monoclonal Antibody, Unconjugated(bsm-60920R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD33 Monoclonal Antibody, Unconjugated(bsm-60920R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.