
MST1/MST2 Rabbit pAb

Catalog Number: bs-8477R

Target Protein: MST1/MST2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ICC/IF (1:50)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Pig, Cow, Dog, Horse)

Predicted MW: 56 kDa

Entrez Gene: 6788

Swiss Prot: Q13043

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene encodes a serine/threonine protein kinase activated by proapoptotic molecules indicating the encoded protein functions as a growth suppressor. Cleavage of the protein product by caspase removes the inhibitory C-terminal portion. The N-terminal portion is transported to the nucleus where it homodimerizes to form the active kinase which promotes the condensation of chromatin during apoptosis. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2012]

PRODUCT SPECIFIC PUBLICATIONS

[IF=5.589] Gao R et al. PM2.5-associated nitro-PAH exposure promotes tumor cell metastasis through Hippo-

YAPmediated transcriptional regulation. Sci Total Environ. 2019 Aug 15;678:611-617. WB ; Human . 31078851

[IF=4.8] Pengfei Li. et al. IL-32 aggravates metabolic disturbance in human nucleus pulposus cells by activating FAT4-mediated Hippo/YAP signaling. INT IMMUNOPHARMACOL. 2024 Nov;141:112966 WB,IP ; Human . 39178518