bs-60174R

[Primary Antibody]

Butyryl-Histone H4 (Lys8) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: Butyryl-Histone H4 (Lys8) **Purification:** Antigen affinity purification

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Histones are basic nuclear proteins that are responsible for the

nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element.

[provided by RefSeq, Jul 2008]


Applications: WB (1:500-2000)

ICC/IF (1:100-500) ChIP (4 ug/5x106 cells)


Reactivity: Human, Mouse, Rat

Subcellular Location: Nucleus

VALIDATION IMAGES



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: (-) HeLa, (+) HeLa+Sodium butyrate (30mM, 4hr) Protein loading quantity: 20 µg Exposure time: 60 s Predicted MW: 11 kDa Observed MW: 11 kDa



Cell line: (A) HeLa, (B) HeLa+SBA (5mM, 24hr)
Fixative: 100% Ice-cold methanol
Permeabilization: 0.1% TritonX-100 Primary ab
dilution: 1:50 Primary incubation condition: 4°C
overnight Secondary ab: Goat Anti-Rabbit IgG
Nuclear counter stain: DAPI (Blue) Counter stain:
Tubulin (Red) Comment: Color green is the
positive signal for bs-60174R