bs-0456R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

GNRH/LHRH Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: GNRH/LHRH

Immunogen: KLH conjugated synthetic peptide ([D-Trp6]-LHRH Fragment, 1-6):

28-40/92.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is secreted and then cleaved to

form the 10 aa luteinizing hormone-releasing hormone (LHRH, also known as gonadoliberin-1), and prolactin release-inhibiting factor (also known as GnRH-associated peptide 1). LHRH stimulates the release of luteinizing and follicle stimulating hormones, which are important for reproduction. Mutation in this gene are associated with hypogonadotropic hypogonadism. Alternatively spliced transcript variants have been described for this gene. [provided by

RefSeq, Jul 2012].

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500)


Reactivity: Human (predicted: Mouse,

Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse)


Predicted MW.: 7.9 kDa

Subcellular Location: Secreted

VALIDATION IMAGES



Tissue/cell: human mammary cancer; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-GNRH Polyclonal Antibody, Unconjugated(bs-0456R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (human kidney tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GNRH) Polyclonal Antibody, Unconjugated (bs-0456R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

- SELECTED CITATIONS -

- [IF=4.57] Liu, Hongxia, et al. "Ultrasound-Mediated Destruction of LHRHa Targeted and Paclitaxel Loaded Lipid Microbubbles Induces Proliferation Inhibition and Apoptosis in Ovarian Cancer Cells." Molecular Pharmaceutics (2013). ICC: Human. 24266423
- [IF=1.514] Liu T et al. Di-(2-ethylhexyl) phthalate induces precocious puberty in adolescent female rats.Iran J Basic Med Sci. 2018 Aug;21(8):848-855. IHC; Rat. 30186573