

bs-11494R**[Primary Antibody]****UNC5D Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 137970	SWISS: Q6UXZ4	IHC-F (1:100-500)
Target: UNC5D		IF (1:100-500)
Immunogen: KLH conjugated synthetic peptide derived from human UNC5H4: 112-200/953. < Extracellular >		ICC/IF (1:100-500)
Purification: affinity purified by Protein A		ELISA (1:5000-10000)
Concentration: 1mg/ml		Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Cow, Chicken, Dog, Horse)
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 102 kDa
Background: The UNC5H family of proteins act as transmembrane receptors for netrin-1 and play a crucial role in axon guidance and migration of neural cells. Additionally, when cleaved by a caspase to produce an intracellular fragment containing a death domain, UNC5H receptors induce apoptosis. This activity is blocked by the binding of netrin-1. In the absence of netrin-1, UNC5H receptors act as tumor suppressors by inhibiting anchorage-independent growth and invasion, but mutation of these receptors provides a potential mechanism for tumorigenicity. The expression of UNC5H receptors is down-regulated in multiple carcinomas, including colorectal, breast, ovary, uterus, stomach, lung, and kidney cancers. UNC5H4, also known as UNC5D (unc-5 homolog D), is single-pass type I membrane protein that is a member of the UNC5H netrin receptor family. Two isoforms of UNC5H4 exist due to alternative splicing events.		Subcellular Location: Cell membrane

— SELECTED CITATIONS —

- **[IF=6.684]** Aditi Karmakar. et al. Identification of Epigenetically Modified Hub Genes and Altered Pathways Associated With Retinoblastoma. Front Cell Dev Biol. 2022; 10: 743224 WB ;Human. 35359459