

**bs-11511R****[ Primary Antibody ]****BBS9 Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**


<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000)
<b>Clonality:</b> Polyclonal		<b>IHC-P</b> (1:100-500)
<b>GeneID:</b> 27241	<b>SWISS:</b> Q3SYG4	<b>IHC-F</b> (1:100-500)
<b>Target:</b> BBS9		<b>IF</b> (1:100-500)
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human BBS9: 244-320/887.		<b>Reactivity:</b> Mouse, Rat (predicted: Human, Rabbit, Pig, Sheep, Dog, Horse)
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		<b>Predicted MW.:</b> 99 kDa
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		<b>Subcellular Location:</b> Cell membrane ,Cytoplasm
<b>Background:</b> BBS9 is an 887 amino acid protein that localizes to both the cytoplasm and the centrosome and exists as six alternatively spliced isoforms. Expressed in a wide variety of tissues, including liver, lung, heart, brain and skeletal muscle, BBS9 functions as a component of the multi-protein BBSome complex which is required for ciliogenesis and is regulated by GDP/GTP exchange factors. Defects in the gene encoding BBS9 are associated with the pathogenesis of Bardet-Biedl syndrome type 9 (BBS9), an autosomal recessive disorder that is characterized by severe pigmentary retinopathy, early onset obesity, polydactyly, hypogenitalism, renal malformation and mental retardation. Additionally, chromosomal aberrations involving the BBS9 gene may play a role in the formation of Wilms tumor 5 (WT5).		

**— VALIDATION IMAGES —**

Sample: Lane 1: Uterus (Mouse) Lysate at 40 ug  
Lane 2: Testis (Rat) Lysate at 40 ug Lane 3: Huvec (Human) Lysate at 30 ug Lane 4: U87MG (Human) Lysate at 30 ug Lane 5: Hela (Human) Lysate at 30 ug Lane 6: 293T (Human) Lysate at 30 ug  
Primary: Anti-BBS9 (bs-11511R) at 1/1000  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 100-110 kD Observed band size: 120 kD



Paraformaldehyde-fixed, paraffin embedded (mouse heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (BBS9) Polyclonal Antibody, Unconjugated (bs-11511R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (BBS9) Polyclonal Antibody, Unconjugated (bs-11511R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.