

bs-18555R

[Primary Antibody]

ZCCHC6 Rabbit pAb

BioSS
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Mouse (predicted: Human, Rat, Rabbit, Pig, Dog) Predicted MW.: 171 kDa Subcellular Location: Cytoplasm ,Nucleus
Clonality: Polyclonal		
GeneID: 79670	SWISS: Q5VYS8	
Target: ZCCHC6		
Immunogen: KLH conjugated synthetic peptide derived from human ZCCHC6: 111-210/1495.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: ZCCHC6 is a 1,495 amino acid uridylyltransferase that mediates RNA uridylation. A member of the DNA polymerase type-B-like family, ZCCHC6 contains three CCHC-type zinc fingers and two PAP-associated domains, and exists as six alternatively spliced isoforms. The gene encoding ZCCHC6 maps to human chromosome 9q21.33 and mouse chromosome 13 B2.		

— VALIDATION IMAGES —



Sample: Liver (Mouse) Lysate at 40 ug Primary:
Anti-ZCCHC6 (bs-18555R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at
1/20000 dilution Predicted band size: 171 kD
Observed band size: 171 kD



Paraformaldehyde-fixed, paraffin embedded
(Mouse brain); Antigen retrieval by boiling in
sodium citrate buffer (pH6.0) for 15min; Block
endogenous peroxidase by 3% hydrogen
peroxide for 20 minutes; Blocking buffer (normal
goat serum) at 37°C for 30min; Antibody
incubation with (ZCCHC6) Polyclonal Antibody,
Unconjugated (bs-18555R) at 1:400 overnight at
4°C, followed by operating according to SP
Kit(Rabbit) (sp-0023) instructions and DAB
staining.