

bs-5348R**[Primary Antibody]****phospho-FXYD1 (Ser83) Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 5348	SWISS: O00168	IHC-F (1:100-500)
Target: FXYD1 (Ser83)		IF (1:100-500)
		ELISA (1:5000-10000)
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human FXYD1 around the phosphorylation site of Ser83: RS(p-S)IR.		Reactivity: (predicted: Human, Mouse, Rat, Pig, Sheep, Cow, Dog)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 10 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cell membrane
Background: PLM (FXYD1) is a member of a family of small membrane proteins that share a 35-amino acid signature sequence domain, beginning with the sequence PFXVD and containing 7 invariant and 6 highly conserved amino acids. FXYD2, also known as the gamma subunit of the Na,K-ATPase, regulates the properties of that enzyme. FXYD1 (phospholemman), FXYD2 (gamma), FXYD3 (MAT-8), FXYD4 (CHIF), and FXYD5 (RIC) have been shown to induce channel activity in experimental expression systems. PLM may be phosphorylated by several kinases, including protein kinase A, protein kinase C, NIMA kinase, and myotonic dystrophy kinase. It is thought to form an ion channel or regulate ion channel activity.		