

bs-5348R**[Primary Antibody]****phospho-FXYD1 (Ser83) Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 5348</p> <p>Target: phospho-FXYD1 (Ser83)</p> <p>Immunogen: KLH conjugated Synthesised phosphopeptide derived from human FXYD1 around the phosphorylation site of Ser83: RS(p-S)IR.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: PLM (FXYD1) is a member of a family of small membrane proteins that share a 35-amino acid signature sequence domain, beginning with the sequence PFXVD and containing 7 invariant and 6 highly conserved amino acids. FXYD2, also known as the gamma subunit of the Na,K-ATPase, regulates the properties of that enzyme. FXYD1 (phospholemman), FXYD2 (gamma), FXYD3 (MAT-8), FXYD4 (CHIF), and FXYD5 (RIC) have been shown to induce channel activity in experimental expression systems. PLM may be phosphorylated by several kinases, including protein kinase A, protein kinase C, NIMA kinase, and myotonic dystrophy kinase. It is thought to form an ion channel or regulate ion channel activity.</p>	<p>Isotype: IgG</p> <p>SWISS: O00168</p> <p>Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000)</p> <p>Reactivity: (predicted: Human, Mouse, Rat, Pig, Sheep, Cow, Dog)</p> <p>Predicted MW.: 10 kDa</p> <p>Subcellular Location: Cell membrane</p>
---	--