

bs-1240G**[Primary Antibody]****human Fibrinogen Goat pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Goat Clonality: Polyclonal GeneID: 2243 Target: human Fibrinogen Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Fibrinogen is the main protein of blood coagulation system. It is a large protein and it consists of two identical subunits that contain three polypeptide chains: alpha, beta and gamma. All chains are connected with each other by a number of disulfide bonds. Fibrinopeptides A (1 to 16 amino acids) and B (1 to 17 amino acids) are released by thrombin from the N terminal parts of alpha and beta chains, respectively. In this way fibrinogen is converted into fibrin, which by means of polymerization forms a fibrin clot. Fibrinogen clotting underlies pathogenesis of MI, thromboembolism and thromboses of arteries and veins, since fibrin is the main substrate for thrombus formation. Fibrinogen activation is also involved in pathogenesis of inflammation, tumor growth and many other diseases. The normal fibrinogen concentration in plasma is about 3 mg/ml. The elevated level of fibrinogen in patient's blood is regarded as an independent risk factor for cardiovascular diseases. An increase in blood fibrinogen concentration was shown to be a strong predictor of coronary heart disease (Sonel A. et al, and Rapold H.J. et al). All these facts make fibrinogen an important parameter in the diagnosis of cardiovascular diseases.	Isotype: IgG SWISS: P02671	Applications: WB (1:500-2000) Reactivity: Human (predicted: Mouse, Rat) Predicted MW.: 240 kDa Subcellular Location: Secreted
---	---	--

— VALIDATION IMAGES —

Sample: Lane 1: Human Plasma Lane 2: Human Fibrinogen (from plasma) (bs-1240P) Primary: Anti-Fibrinogen (bs-1240G) at 1/1000 dilution Secondary: IRDye800CW Donkey anti-Goat IgG at 1/20000 dilution Predicted band size: 240 kDa Observed band size: 55,65,73 kDa

— SELECTED CITATIONS —

- **[IF=4.882]** Li Peichuang, et al. Preparation of phospholipid-based polycarbonate urethanes for potential applications of blood-contacting implants. Regen Biomater. 2020 Oct;7(5):491-504 ELISA ;Rabbit. 33149938