

bs-0267R**[Primary Antibody]****PRKCB Rabbit pAb**

Bioss
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit Clonality: Polyclonal GeneID: 5579 Target: PRKCB Immunogen: KLH conjugated synthetic peptide derived from human PKC beta 1: 601-673/673. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This protein kinase has been reported to be involved in many different cellular functions, such as B cell activation, apoptosis induction, endothelial cell proliferation, and intestinal sugar absorption. Studies in mice also suggest that this kinase may also regulate neuronal functions and correlate fear-induced conflict behavior after stress. Alternatively spliced transcript variants encoding distinct isoforms have been reported.	Isotype: IgG SWISS: P05771	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1:ug/Test) Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Cow) Predicted MW.: 74 kDa Subcellular Location: Cell membrane ,Cytoplasm ,Nucleus
---	---	--

— VALIDATION IMAGES —


Sample: spleen (Mouse) Lysate at 40 ug Primary:
 Anti-PKC beta 1 + 2 (bs-0267R) at 1/300 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at
 1/20000 dilution Predicted band size: 74 kD
 Observed band size: 74 kD



Tissue/cell: human kidney tissue; 4%
 Paraformaldehyde-fixed and paraffin-
 embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block
 endogenous peroxidase by 3% Hydrogen
 peroxide for 30min; Blocking buffer (normal goat
 serum,C-0005) at 37°C for 20 min; Incubation:
 Anti- PKC beta-1+2 Polyclonal Antibody,
 Unconjugated(bs-0267R) 1:200, overnight at 4°C,
 followed by conjugation to the secondary
 antibody(SP-0023) and DAB(C-0010) staining



Blank control (Black line): Molt4 (Black). Primary
 Antibody (green line):Rabbit Anti-PKC beta 1 + 2
 antibody (bs-0267R) Dilution:1µg /10⁶ cells;
 Isotype Control Antibody (orange line): Rabbit
 IgG . Secondary Antibody (white blue line): Goat
 anti-rabbit IgG-AF647 Dilution: 1µg /test.
 Protocol The cells were fixed with 4% PFA
 (10min at room temperature)and then
 permeabilized with 90% ice-cold methanol for
 20 min at room temperature. The cells were then
 incubated in 5%BSA to block non-specific
 protein-protein interactions for 30 min at room
 temperature .Cells stained with Primary
 Antibody for 30 min at room temperature. The
 secondary antibody used for 40 min at room
 temperature. Acquisition of 20,000 events was
 performed.



Blank control: Molt4. Primary Antibody (green line): Rabbit Anti-PKC beta 1 + 2 antibody (bs-0267R) Dilution: 1µg /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=3.69]** Ji-Sheng Wang. et al. Effect of leech-centipede medicine on improving erectile function in DIED rats via PKC signalling pathway-related molecules. J Ethnopharmacol. 2021 Mar;267:113463 WB ;Rat. 33049347
- **[IF=4.36]** Xiao Li. et al. Mechanism by which Huoxue Tongluo Qiwei Decoction improves the erectile function of rats with diabetic erectile dysfunction. J Ethnopharmacol. 2022 Jan;283:114674 WB ;Rat. 10.1016/j.jep.2021.114674
- **[IF=2]** Penglei Ge. et al. Identifying drug candidates for pancreatic ductal adenocarcinoma based on integrative multiomics analysis. J GASTROINTEST ONCOL. 2024 Jun 30; 15(3): 1265–1281 IHC,WB ;Human. 38989421