

---

## PP-1B Mouse pAb

Catalog Number: bs-0133M

Target Protein: PP-1B

Concentration: 1mg/ml

Form: Liquid

Host: Mouse

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Rat (predicted:Human, Mouse, Rabbit, Cow, Horse)

Predicted MW: 37 kDa

Subcellular Nucleus

Locations:

Entrez Gene: 5500

Swiss Prot: P62140

Source: KLH conjugated synthetic peptide derived from human PP-1B: 155-250/327.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** The phosphorylation and de phosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions in eukaryotes, including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the serine/ threonine protein phosphatases. The protein phosphatases (PP) holoenzyme is a trimetric complex composed of a regulatory subunit, a variable subunit and a catalytic subunit. Four major families of protein phosphatase catalytic subunits have been identified and are designated PP1, PP2A, PP2B (calcineurin) and PP2C. An additional protein phosphatase catalytic subunit, PPX or PP4, is the member of a potentially novel PP family. The PP1 family is composed of subfamily members PP1 alpha, PP1 beta, and PP1 gamma. The PP2A family is composed of subfamily members PP2A alpha, and PP2A beta. The PP2B family is composed of subfamily members PP2B-A alpha, PP2B-A beta and PP2B-A gamma. The PP2C family is composed of PP2C alpha, PP2C beta, PP2C gamma. Wip1, a protein identified in the p53 DNA response pathway, has also been identified as a potential member of the PP2C family. Regulatory subunits include nuclear inhibitor of PP1 (NIPP1), PP2A-A1 alpha and -A beta, PP2A-B alpha and B beta, PP2A-C alpha and C beta, PP2A-B56-

alpha and-B56 beta, PR48 and PP2B-B1 and-B2.

## VALIDATION IMAGES

---



Tissue/cell: rat pancreas tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-PP-1B Polyclonal Antibody, Unconjugated(bs-0133M) 1:300, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0024) and DAB(C-0010) staining