

bs-6742R

[**Primary Antibody**]

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

TPD52 Rabbit pAb

— DATASHEET —

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 7163</p> <p>Target: TPD52</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human TPD52: 141-190/224.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: Isoform 2 is expressed in colon, breast, prostate, pancreas and kidney tumor cell lines. Isoform 2 is expressed at high levels in kidney, prostate, brain, small intestine and pancreas, at moderate levels in placenta and colon, at low levels in lung, liver and heart, and at very low levels in spleen, thymus, peripheral mononuclear blood cells, testis and ovary.</p>	<p>Isotype: IgG</p> <p>SWISS: P55327</p>	<p>Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000)</p> <p>Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Sheep, Cow)</p> <p>Predicted MW.: 24 kDa</p> <p>Subcellular Location: Cytoplasm</p>
---	--	--

— SELECTED CITATIONS —

- **[IF=5.17]** Wang et al. Decreased TPD52 expression is associated with poor prognosis in primary hepatocellular carcinoma. (2016) Oncotarget. 7:6323-34 IHC ;Human. 26575170