

bs-6749R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Ubiquitously expressed transcript Rabbit pAb**— DATASHEET —**

| | | |
|---|---|--|
| Host: Rabbit Clonality: Polyclonal GeneID: 8409 Target: Ubiquitously expressed transcript Immunogen: KLH conjugated synthetic peptide derived from human UXT/Ubiquitously expressed transcript: 85-157/157. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: UXT is a novel protein, highly conserved in mouse. It interacts with the N terminus of the androgen receptor and plays a role in facilitating receptor-induced transcriptional activation. It is abundantly expressed in tumor tissues and is likely to be involved in tumorigenesis. | Isotype: IgG SWISS: Q9UBK9 | Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse, Rat (predicted: Pig, Sheep, Cow, Dog, Horse) Predicted MW.: 18 kDa Subcellular Location: Cytoplasm ,Nucleus |
|---|---|--|

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Mouse stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Ubiquitously expressed transcript) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Protein UXT) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

— SELECTED CITATIONS —

- **[IF=5.17]** Tew et al. Vitamin K epoxide reductase regulation of androgen receptor activity. (2017) Oncotarget. 8:13818-13831 WB ;Homo sapiens (Human). 28099154