

bs-60035R

[Primary Antibody]


Acetyl-Histone H4 (Lys12) Rabbit pAb



- VALIDATION IMAGES



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: (-) HeLa, (+) HeLa+Sodium butyrate (30mM, 4hr) Protein loading quantity: 20 µg Exposure time: 60 s Predicted MW: 11 kDa Observed MW: 11 kDa



Tissue: Human neuroblastoma Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:200 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit) (sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bs-60035R

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

Applications: WB (1:500-1:2000) **IHC-P** (1:100-500) IHC-F (1:100-500) ICC/IF (1:50-1:200)

Reactivity: Human (predicted: Mouse,