## bsm-60574R

# [ Primary Antibody ]

# BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## **ELP3 Recombinant Rabbit mAb**

- DATASHEET -

Host: Rabbit Isotype: IgG/Kappa
Clonality: Recombinant CloneNo.: E9G9

Target: ELP3

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** ELP3 is the catalytic subunit of the histone acetyltransferase

elongator complex, which contributes to transcript elongation and also regulates the maturation of projection neurons (Creppe et al.,  $\frac{1}{2} = \frac{1}{2} \left( \frac{1}{2} - \frac{1}{2} \right) \left( \frac{1}{2} - \frac{1}{2} - \frac{1}{2} \right) \left( \frac{1}{2} - \frac{1}{2} - \frac{1}{2} - \frac{1}{2} \right) \left( \frac{1}{2} - \frac{1}{$ 

2009 [PubMed 19185337]).[supplied by OMIM, Apr 2009]

**Applications: WB** (1:500-1:1000)

**IHC-P** (1:50-1:100)

IHC-F

ICC/IF (1:50)


**Reactivity:** Human, Mouse

(predicted: Rat)


Predicted MW.: 62 kDa

**Subcellular Location:** Cell membrane , Nucleus


### VALIDATION IMAGES




Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:1000 Primary ab incubation condition: 2 hours at room temperature
Secondary ab: Goat Anti-Rabbit IgG H&L (HRP)
Lysate: HeLa, K562, Mouse brain, Raw264.7
Protein loading quantity: 20 µg Exposure time:
60 s Predicted MW: 62 kDa Observed MW: 62 kDa




Tissue: Human tonsil Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit) (sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60574R



Tissue: Mouse testis Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100 Primary ab incubation condition: 1 hour at room temperature Secondary ab:SP Kit(Rabbit) (sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60574R



Cell line: HeLa Fixative: 100% Ice-cold methanol Permeabilization: 0.1% TritonX-100 Primary ab dilution: 1:50 Primary incubation condition: 4°C overnight Secondary ab: Goat Anti-Rabbit IgG Nuclear counter stain: DAPI (Blue) Comment: Color green is the positive signal for bsm-60574R



1