bs-2428R

[Primary Antibody]

IL-9 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –––––		400-901-9800
Host: Rabbit Clonality: Polyclonal	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500)
GenelD: 16198	SWISS: P15247	IF (1:200-800)
Target: IL-9 Immunogen: KLH conjugated synthetic peptide derived from mouse IL-9: 21-120/144.		Reactivity: (predicted: Mouse, Rat)
Purification: affinity purified by Protein A Concentration: 1mg/ml		Predicted MW.: ^{16 kDa}
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Secreted
Background: The protein encoded by this gene is a cytokine that actsas a regulator of a variety of hematopoietic cells. This cytokinestimulates cell proliferation and prevents apoptosis. It functionsthrough the interleukin 9 receptor (IL9R), which activatesdifferent signal transducer and activator (STAT) proteins and thusconnects this cytokine to various biological processes. The geneencoding this cytokine has been identified as a candidate gene forasthma. Genetic studies on a mouse model of asthma demonstratedthat this cytokine is a determining factor in the pathogenesis ofbronchial hyperresponsiveness. [provided by RefSeq, Jul 2008].		

- SELECTED CITATIONS -----

- [IF=5.6] Yun Sun. et al. Inhibitory effects of catalpol on DNCB-induced atopic dermatitis and IgE-mediated mast cells reaction. INT IMMUNOPHARMACOL. 2024 Jan;126:111274 ELISA, IHC ;MOUSE. 38041954
- [IF=3.7] Yao, Xiuhua, et al. "Neutralization of IL 9 ameliorates symptoms of experimental autoimmune myasthenia gravis in rats by decreasing effector T cells and altering humoral responses." Immunology (2014). Other ;="Rat". 24850614
- [IF=1.5] Lin, Yinyao, et al. "Expression of interleukin-9 and its upstream stimulating factors in rats with ischemic stroke." Neurological Sciences (2015): 1-8. IHC ;="Rat". 25652434