

**bs-3768R****[ Primary Antibody ]****Histone H3 (tri methyl K36) Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**


<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>Flow-Cyt</b> (1µg /Test)
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 8350	<b>SWISS:</b> P68431	
<b>Target:</b> Histone H3 (tri methyl K36)		
<b>Immunogen:</b> KLH conjugated synthesised methylpeptide derived from human Histone H3 around the methylation site of tri methyl K36: GV(Tri Methyl-K)KP.		
<b>Purification:</b> affinity purified by Protein A		<b>Reactivity:</b> Human, Mouse, Rat (predicted: Rabbit, Pig, Cow, Fruit Fly)
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> Modulation of the chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of four core histone proteins (H2A, H2B, H3 and H4), is the primary building block of chromatin. The N-terminal tail of core histones undergoes different posttranslational modifications including acetylation, phosphorylation and methylation. These modifications occur in response to cell signal stimuli and have a direct effect on gene expression. In most species, the histone H2B is primarily acetylated at lysines 5, 12, 15 and 20. Histone H3 is primarily acetylated at lysines 9, 14, 18 and 23. Acetylation at lysine 9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms. Phosphorylation at Ser10 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis.		
		<b>Predicted MW.:</b> 15 kDa
		<b>Subcellular Location:</b> Nucleus

**— VALIDATION IMAGES —**


Sample: Lane 1: SiHa (Human) Cell Lysate at 30 ug  
 Lane 2: MCF-7 (Human) Cell Lysate at 30 ug  
 Lane 3: U-2OS (Human) Cell Lysate at 30 ug  
 Lane 4: Jurkat (Human) Cell Lysate at 30 ug  
 Primary: Anti-Histone H3 (tri methyl K36) (bs-3768R) at 1/1000 dilution  
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
 Predicted band size: 17 kD  
 Observed band size: 17 kD



Paraformaldehyde-fixed, paraffin embedded (mouse pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Histone H3 (tri methyl K36)) Polyclonal Antibody, Unconjugated (bs-3768R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: rat kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Histone H3 (tri methyl K36) Polyclonal Antibody, Unconjugated (bs-3768R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



Blank control (blue line): Hela (fixed with 70% ethanol (Overnight at 4°C) and then permeabilized with 90% ice-cold methanol for 30 min on ice). Primary Antibody (green line): Rabbit Anti-Histone H3 (tri methyl K36) antibody (bs-3768R), Dilution: 1µg /10<sup>6</sup> cells; Isotype Control Antibody (orange line): Rabbit IgG . , Dilution: 1µg /test.

## — SELECTED CITATIONS —

- **[IF=4.907]** Shen Y et al. Set7 Is a H3K37 Methyltransferase in Schizosaccharomyces pombe and Is Required for Proper Gametogenesis. Structure. 2019 Feb 12. pii: S0969-2126(19)30011-5. WB ;Yeast(S. pombe). 30773398
- **[IF=4]** Daoyu Zhang. et al. Transcription factor TFAP2C affects porcine early embryo development via regulating epigenetic modification. REPROD BIOMED ONLINE. 2023 Dec;:103772 IF ;Pig. 10.1016/j.rbmo.2023.103772
- **[IF=2.626]** Zhao, Gang. et al. LINC02381, a sponge of miR-21, weakens osteogenic differentiation of hUC-MSCs through KLF12-mediated Wnt4 transcriptional repression. 2021 Nov 15 WB ;Mouse,Human. 34778905