

bs-20711R**[Primary Antibody]****B7H4 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Sheep, Cow)
GeneID: 79679	SWISS: Q7Z7D3	Predicted MW.: 28 kDa
Target: B7H4		Subcellular Location: Cell membrane
Immunogen: KLH conjugated synthetic peptide derived from human B7H4: 151-250/282. < Extracellular >		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: B7-H4 protein is expressed on the surface of a variety of immune cells and functions as a negative regulator of T cell responses. While B7-H4 mRNA is widely distributed in mouse and human peripheral tissues, cell surface expression of B7-H4 protein is limited and shows an inducible pattern on hematopoietic cells. Putative receptor of B7-H4 can be upregulated on activated T cells. By arresting cell cycle, B7-H4 ligation of T cells has a profound inhibitory effect on the growth, cytokine secretion, and development of cytotoxicity. Administration of B7-H4Ig into mice impairs antigen-specific T cell responses whereas blockade of endogenous B7-H4 by specific monoclonal antibody promotes T cell responses. B7-H4 thus may participate in negative regulation of cell-mediated immunity in peripheral tissues.		