

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## ATG14 Rabbit pAb

Catalog Number: bs-7462R

Target Protein: ATG14
Concentration: 1mg/ml

Form: Liquid

Host: Rabbit
Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human (predicted: Mouse, Rat, Pig, Cow, Dog)

Predicted MW: 55 kDa

Subcellular Cytoplasm

Locations:

Entrez Gene: 22863 Swiss Prot: Q6ZNE5

Source: KLH conjugated synthetic peptide derived from human ATG14: 41-140/492.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: Required for both basal and inducible autophagy. Plays a role in autophagosome formation

and MAP1LC3/LC3 conjugation to phosphatidylethanolamine. Promotes BECN1

translocation from the trans-Golgi network to autophagosomes. Enhances PIK3C3 activity in

a BECN1-dependent manner.

## **VALIDATION IMAGES**



Sample: Lane 1: Human MCF-7 cell lysates Lane 2: Human SH-SY5Y cell lysates Lane 3: Human HeLa cell lysates Lane 4: Human A549 cell lysates Lane 5: Human A431 cell lysates Primary: Anti-ATG14 (bs-7462R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 55 kDa Observed band size: 68 kDa



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ATG14) Polyclonal Antibody, Unconjugated (bs-7462R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

## PRODUCT SPECIFIC PUBLICATIONS

[IF=9.584] Zheng, Bingxin. et al. Siglec-15-induced autophagy promotes invasion and metastasis of human osteosarcoma cells by activating the epithelial–mesenchymal transition and Beclin-1/ATG14 pathway. CELL BIOSCI. 2022 Dec;12(1):1-15 WB; Human. 35842729