

bs-7462R**[Primary Antibody]****ATG14 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 22863 Target: ATG14 Immunogen: KLH conjugated synthetic peptide derived from human ATG14: 41-140/492. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Required for both basal and inducible autophagy. Plays a role in autophagosome formation and MAP1LC3/LC3 conjugation to phosphatidylethanolamine. Promotes BECN1 translocation from the trans-Golgi network to autophagosomes. Enhances PIK3C3 activity in a BECN1-dependent manner.	Isotype: IgG SWISS: Q6ZNE5	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human (predicted: Mouse, Rat, Pig, Cow, Dog) Predicted MW.: 55 kDa Subcellular Location: Cytoplasm
---	---	---

— VALIDATION IMAGES —

Sample: Lane 1: Human MCF-7 cell lysates Lane 2: Human SH-SY5Y cell lysates Lane 3: Human HeLa cell lysates Lane 4: Human A549 cell lysates Lane 5: Human A431 cell lysates Primary: Anti-ATG14 (bs-7462R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 55 kDa Observed band size: 68 kDa



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ATG14) Polyclonal Antibody, Unconjugated (bs-7462R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=9.584]** Zheng, Bingxin. et al. Siglec-15-induced autophagy promotes invasion and metastasis of human osteosarcoma cells by activating the epithelial-mesenchymal transition and Beclin-1/ATG14 pathway. CELL BIOSCI. 2022 Dec;12(1):1-15 WB ;Human. 35842729