bs-11897R

[Primary Antibody]


NEURL Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -Host: Rabbit Isotype: IgG Applications: IHC-P (1:100-500) IHC-F (1:100-500) Clonality: Polyclonal IF (1:100-500) GenelD: 9148 SWISS: 076050 Target: NEURL Immunogen: KLH conjugated synthetic peptide derived from human NEURL/RNF67: 51-150/574. Purification: affinity purified by Protein A Reactivity: Mouse, Rat Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated Predicted MW.: ^{62 kDa} freeze/thaw cycles. Background: NEURL is involved in the determination of cell fate in the neurogenic region of the embryo and plays a role in the

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (rat pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (NEURL) Polyclonal Antibody, Unconjugated (bs-11897R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



determination of cell fate in the central nervous system. NEURL may act as a tumor suppressor whose inactivation could be associated with malignant progression of astrocytic tumors.

> Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NEURL) Polyclonal Antibody, Unconjugated (bs-11897R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

(predicted: Human, Pig, Sheep, Cow, Chicken, Dog)

Subcellular Cell membrane ,Cytoplasm