

**bs-5999R****[ Primary Antibody ]****DAB2 Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**


<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500)
<b>Clonality:</b> Polyclonal		<b>IHC-F</b> (1:100-500)
<b>GeneID:</b> 1601	<b>SWISS:</b> P98082	<b>IF</b> (1:100-500)
<b>Target:</b> DAB2		<b>Flow-Cyt</b> (1µg/Test)
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human DAB2: 31-130/770.		<b>Reactivity:</b> Human, Mouse (predicted: Rat, Rabbit, Pig, Cow, Chicken, Dog, GuineaPig, Horse)
<b>Purification:</b> affinity purified by Protein A		<b>Predicted MW.:</b> 82 kDa
<b>Concentration:</b> 1mg/ml		<b>Subcellular Location:</b> Cytoplasm
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> This gene encodes a mitogen-responsive phosphoprotein. It is expressed in normal ovarian epithelial cells, but is down-regulated or absent from ovarian carcinoma cell lines, suggesting its role as a tumor suppressor. This protein binds to the SH3 domains of GRB2, an adaptor protein that couples tyrosine kinase receptors to SOS (a guanine nucleotide exchange factor for Ras), via its C-terminal proline-rich sequences, and may thus modulate growth factor/Ras pathways by competing with SOS for binding to GRB2. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2011].		

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (DAB2) Polyclonal Antibody, Unconjugated (bs-5999R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human adrenal gland); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (DAB2) Polyclonal Antibody, Unconjugated (bs-5999R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: HeLa. Primary Antibody (green line): Rabbit Anti-DAB2 antibody (bs-5999R) Dilution: 1µg / 10<sup>6</sup> cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat anti-rabbit IgG-AF647 Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

**— SELECTED CITATIONS —**

- **[IF=2.751]** Chun-Lian Liang, et al. DAB2 promotes pulmonary fibrosis and may act as an intermediate between IGF-1R and PI3K/AKT signaling pathways. EXP THER MED. 2023 Apr;25(4):1-11 WB ;Human. 37021069

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

- **[IF=2.127]** Zhou J et al. Phosphorylation of Dab2 is involved in inhibited VEGF - VEGFR - 2 signaling induced by downregulation of Syndecan - 1 in glomerular endothelial cell. Cell Biology International. WB ;Rat. doi:10.1002/cbin.11288