bs-6005R

[Primary Antibody]

EXTL2 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Rabbi	t Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyc	lonal	IHC-F (1:100-500) IF (1:100-500)
GenelD: 2135	SWISS: Q9UBQ6	ELISA (1:5000-10000)
Target: EXTL2	2	Reactivity: (predicted: Human, Mouse,
Immunogen: KLH conjugated synthetic peptide derived from human EXTL: 5-110/330.		
Purification: affinit	y purified by Protein A	
Concentration: 1mg/ml		Predicted MW.: ^{37 kDa}
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		50% Subcellular Secreted ,Cell membrane
hepar beta-	2 is a glycosyltransferase required for the biosynth an-sulfate and is responsible for the alternating a 1-4-linked glucuronic acid (GlcA) and alpha-1-4-lin glucosamine (GlcNAc) units to nascent heparan s s.	ddition of nked N-

- SELECTED CITATIONS -------

• [IF=3.8] Zhaorui Wang. et al. Metabolomics- and Proteomics-Based Disease Diagnostic Classifier Model for the

Prediction and Diagnosis of Colorectal Carcinoma..JOURNAL OF PROTEOME RESEARCH.2025 Mar 19. IHC ;Human. 40108892