

bs-6005R

[Primary Antibody]

EXTL2 Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000)
Clonality: Polyclonal		
GeneID: 2135	SWISS: Q9UBQ6	
Target: EXTL2		
Immunogen: KLH conjugated synthetic peptide derived from human EXTL: 5-110/330.		
Purification: affinity purified by Protein A		Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Sheep, Dog, Horse)
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 37 kDa
Background: EXTL2 is a glycosyltransferase required for the biosynthesis of heparan-sulfate and is responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA) and alpha-1-4-linked N-acetylglucosamine (GlcNAc) units to nascent heparan sulfate chains.		Subcellular Location: Secreted ,Cell membrane ,Cytoplasm

— SELECTED CITATIONS —

- **[IF=3.8]** Zhaorui Wang. et al.Metabolomics- and Proteomics-Based Disease Diagnostic Classifier Model for the Prediction and Diagnosis of Colorectal Carcinoma..JOURNAL OF PROTEOME RESEARCH.2025 Mar 19. IHC ;Human. 40108892