

bs-0111R**[Primary Antibody]**

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

progesterone receptor Rabbit pAb


DATASHEET

Host: Rabbit Clonality: Polyclonal Target: progesterone receptor Immunogen: KLH conjugated synthetic peptide derived from human progesterone receptor: 501-600/933. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	Isotype: IgG GenelD: 5241	Applications: WB (1:500-2000) Flow-Cyt (1ug/Test)
Background: Estrogen and progesterone receptor are members of a family of transcription factors that are regulated by the binding of their cognate ligands. The interaction of hormone-bound estrogen receptors with estrogen responsive elements(EREs) alters transcription of ERE-containing genes. The carboxy terminal region of the estrogen receptor contains the ligand binding domain, the amino terminus serves as the transactivation domain, and the DNA binding domain is centrally located. Two forms of estrogen receptor have been identified, ER alpha and ER beta. ER alpha and ER beta have been shown to be differentially activated by various ligands. The biological response to progesterone is mediated by two distinct forms of the human progesterone receptor (hPR-A and hPR-B), which arise from alternative splicing. In most cells, hPR-B functions as a transcriptional activator of progesterone-responsive gene, whereas hPR-A function as a transcriptional inhibitor of all steroid hormone receptors.		Reactivity: Human, Rat (predicted: Mouse, Rabbit, Pig, Dog, Horse) Predicted MW: 103 kDa Subcellular Location: Cytoplasm ,Nucleus


VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with progesterone receptor polyclonal antibody, unconjugated (bs-0111R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Blank control:K562. Primary Antibody (green line): Rabbit Anti-progesterone receptor antibody (bs-0111R) Dilution: 2μg /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-PE Dilution: 1μg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control: MCF7. Primary Antibody (green line): Rabbit Anti-progesterone receptor antibody (bs-0111R) Dilution: 1μg /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1μg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=5.546]** Xiangguo Wang, et al. Exosomes from bovine endometrial epithelial cells ensure trophoblast cell development by miR - 218 targeting secreted frizzled related protein 2. 2020 Nov 23 WB ;Bovine. 33230823
- **[IF=3.86]** Chu, Meiqiang, et al. "MicroRNA-126 participates in lipid metabolism in mammary epithelial cells." Molecular and Cellular Endocrinology (2017). WB ;="Human". 28599789
- **[IF=4.1]** Derya Erisik, et al. Differences and Similarities between Colorectal Cancer Cells and Colorectal Cancer Stem Cells: Molecular Insights and Implications. ACS OMEGA. 2023;XXXX(XXX):XXX-XXX ICC ;Human. 37636966