
presenilin 1 Rabbit pAb

Catalog Number: bs-0024R

Target Protein: presenilin 1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), Flow-Cyt (1ug/test), ICC/IF (1:100)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Pig, Cow, Dog)

Predicted MW: 34/52 kDa

Entrez Gene: 5663

Swiss Prot: P49768

Source: KLH conjugated synthetic peptide derived from human Presenilin-1 NTF subunit: 10-80/467.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Alzheimer's disease (AD) patients with an inherited form of the disease carry mutations in the presenilin proteins (PSEN1; PSEN2) or in the amyloid precursor protein (APP). These disease-linked mutations result in increased production of the longer form of amyloid-beta (main component of amyloid deposits found in AD brains). Presenilins are postulated to regulate APP processing through their effects on gamma-secretase, an enzyme that cleaves APP. Also, it is thought that the presenilins are involved in the cleavage of the Notch receptor, such that they either directly regulate gamma-secretase activity or themselves are protease enzymes. Several alternatively spliced transcript variants encoding different isoforms have been identified for this gene, the full-length nature of only some have been determined. [provided by RefSeq, Aug 2008]


VALIDATION IMAGES




Sample: K562 Cell (Human) Lysate at 40 ug HepG2 Cell (Human) Lysate at 40 ug Primary: Anti-presenilin 1 (bs-0024R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34/52 kD Observed band size: 50 kD




Sample: K562 Cell Lysate at 40 ug Primary: Anti- presenilin1 (bs-0024R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34/52 kD Observed band size: 48 kD




Sample: Lane 1: Mouse Brain Lysates Lane 2: Mouse Spinal cord Lysates Lane 3: Mouse Testis Lysates Lane 4: Rat Spinal cord Lysates Lane 5: Rat Testis Lysates Lane 6: Human U251 cell Lysates Lane 7: Human SH-SY5Y cell Lysates Lane 8: Human MCF-7 cell Lysates Lane 9: Human THP-1 cell Lysates Lane 10: Human U-87 MG cell Lysates Primary: Anti-presenilin 1 (bs-0024R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34/52kDa Observed band size: 50kDa



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (presenilin 1) Polyclonal Antibody, Unconjugated (bs-0024R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: Raji. Primary Antibody (green line): Rabbit Anti-presenilin 1 antibody (bs-0024R) Dilution: 1μg /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-PE Dilution: 1μg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



SH-SY5Y cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (presenilin 1) polyclonal Antibody, Unconjugated (bs-0024R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

PRODUCT SPECIFIC PUBLICATIONS

[IF=3.58] Wang et al. YXQN Reduces Alzheimer's Disease-Like Pathology and Cognitive Decline in APPswePS1dE9 Transgenic Mice. (2017) Front.Aging.Neurosci. 9:157 Other ; Mouse . 28603494

[IF=3.457] G V et al. Therapeutic impact of rHuEPO on abnormal platelet APP, BACE 1, presenilin 1, ADAM 10 and Aβ expressions in chronic kidney disease patients with cognitive dysfunction like Alzheimer's disease: A pilot study.Biomed Pharmacother. 2018 Aug;104:211-222. WB ; Human . 29775888

[IF=2.454] Yang Q et al. Ginsenoside Compound K Regulates Amyloid β via the Nrf2/Keap1 Signaling Pathway in Mice with Scopolamine

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Hydrobromide-Induced Memory Impairments.(2018) J Mol Neurosci. WB ; Mouse . 30535776