bs-1221R

[Primary Antibody]

www.bioss.com.cn

sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Rad51 Rabbit pAb

GeneID: 5888 SWISS: Q06609

Target: Rad51

Immunogen: KLH conjugated synthetic peptide derived from human Rad51:

241-339/339.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is a member of the RAD51

utilizing alternative polyA signals exist.

protein family. RAD51 family members are highly similar to bacterial RecA and Saccharomyces cerevisiae Rad51, and are known to be involved in the homologous recombination and repair of DNA. This protein can interact with the ssDNA-binding protein RPA and RAD52, and it is thought to play roles in homologous pairing and strand transfer of DNA. This protein is also found to interact with BRCA1 and BRCA2, which may be important for the cellular response to DNA damage. BRCA2 is shown to regulate both the intracellular localization and DNA-binding ability of this protein. Loss of these controls following BRCA2 inactivation may be a key event leading to genomic instability and tumorigenesis. Two alternatively spliced transcript variants of this gene, which encode distinct proteins, have been reported. Transcript variants

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (1ug/Test) ICC/IF (1:25)

Reactivity: Human, Mouse

(predicted: Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW.: 37 kDa

Subcellular Cytoplasm ,Nucleus

— SELECTED CITATIONS —

• [IF=2.4] Qianwen Chang, et al. Elevated temperature affects the expression of signaling molecules in quail testes meiosis I prophase, but spermatogenesis remains normal. THERIOGENOLOGY. 2024 Aug;: IHC, WB; Quail. 39142066