bs-0516R

[Primary Antibody]

RARB Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Isotype: IgG SWISS: P10826 ic peptide derived from human RARB: in A .% BSA, 0.02% Proclin300 and 50% -20°C for one year. Avoid repeated	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1ug/Test) ICC/IF (1:100) Reactivity: Human, Mouse (predicted: Rat, Pig, Cow, Chicken, Dog) Predicted MW.: 50 kDa
SWISS: P10826 ic peptide derived from human RARB: in A .% BSA, 0.02% Proclin300 and 50% -20°C for one year. Avoid repeated	IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1ug/Test) ICC/IF (1:100) Reactivity: Human, Mouse (predicted: Rat, Pig, Cow, Chicken, Dog) Predicted MW.: ⁵⁰ kDa
SWISS: P10826 ic peptide derived from human RARB: in A .% BSA, 0.02% Proclin300 and 50% -20°C for one year. Avoid repeated	IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1ug/Test) ICC/IF (1:100) Reactivity: Human, Mouse (predicted: Rat, Pig, Cow, Chicken, Dog) Predicted MW.: ⁵⁰ kDa
ic peptide derived from human RARB: in A .% BSA, 0.02% Proclin300 and 50% -20°C for one year. Avoid repeated	Flow-Cyt (1ug/Test) ICC/IF (1:100) Reactivity: Human, Mouse (predicted: Rat, Pig, Cow, Chicken, Dog) Predicted MW.: ⁵⁰ kDa
ic peptide derived from human RARB: in A .% BSA, 0.02% Proclin300 and 50% -20°C for one year. Avoid repeated	Reactivity: Human, Mouse (predicted: Rat, Pig, Cow, Chicken, Dog) Predicted MW.: ⁵⁰ kDa
in A .% BSA, 0.02% Proclin300 and 50% -20°C for one year. Avoid repeated	(predicted: Rat, Pig, Cow, Chicken, Dog) Predicted MW.: ⁵⁰ kDa
.% BSA, 0.02% Proclin300 and 50% -20°C for one year. Avoid repeated	Predicted MW.: ^{50 kDa}
% BSA, 0.02% Proclin300 and 50% -20°C for one year. Avoid repeated	Predicted MW.: ^{50 kDa}
es of vitamin A (retinal) and are believed to haling molecules during vertebrate differentiation. Two families of retinoid ntified. Retinoic acid receptors (RARs) Beta and RAR gamma, each of which has a retinoic acids and belongs to the same otion factors as thyroid hormone ceptor and ecdysone receptor. The ligand RARs are highly conserved and RAR n distinct patterns throughout mature organism. Members of the retinoic RXR alpha, RXR Beta and RXR gamma, are c acid, a stereo and photoisomer of all	Subcellular Location: Cytoplasm ,Nucleus
	is of vitamin A (retinal) and are believed to ialing molecules during vertebrate differentiation. Two families of retinoid ntified. Retinoic acid receptors (RARs) 3eta and RAR gamma, each of which has a retinoic acids and belongs to the same otion factors as thyroid hormone ceptor and ecdysone receptor. The ligand XARs are highly conserved and RAR n distinct patterns throughout mature organism. Members of the retinoic RXR alpha, RXR Beta and RXR gamma, are c acid, a stereo and photoisomer of all d in vivo in both liver and kidney and may hormone.

Sample: Cerebrum (Mouse) Lysate at 40 ug Primary: Anti-RAR Beta (bs-0516R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 50 kD Observed band size: 50 kD



Sample: Lane 1: Mouse Eye tissue lysates Lane 2: Mouse Cerebrum tissue lysates Primary: Anti-RAR Beta (bs-0516R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti- Rabbit IgG at 1/20000 dilution Predicted band size: 50 kDa Observed band size: 50 kDa



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RAR Beta) Polyclonal Antibody, Unconjugated (bs-0516R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.







Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RAR Beta) Polyclonal Antibody, Unconjugated (bs-0516R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining. U87MG cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (RAR Beta) polyclonal Antibody, Unconjugated (bs-0516R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei. Blank control: SHSY5Y. Primary Antibody (green line): Rabbit Anti-RAR Beta antibody (bs-0516R) Dilution: 1ug/Test; Secondary Antibody : Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature.