

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Resistin Rabbit pAb

Catalog Number: bs-0795R

Target Protein: Resistin Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ELISA (1:5000-10000)

Reactivity: Rat (predicted:Mouse)

Predicted MW: 12 kDa
Entrez Gene: 56729
Swiss Prot: Q9HD89

Source: KLH conjugated synthetic peptide derived from human Resistin: 21-114/114.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: This gene belongs to the family defined by the mouse resistin-like genes. The characteristic

feature of this family is the C-terminal stretch of 10 cys residues with identical spacing. The

mouse homolog of this protein is secreted by adipocytes, and may be the hormone

potentially linking obesity to type II diabetes. [provided by RefSeq]

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (rat breast); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Resistin) Polyclonal Antibody, Unconjugated (bs-0795R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

PRODUCT SPECIFIC PUBLICATIONS

[IF=7.419] Hirona Kugo. et al. Tricaprin can prevent the development of AAA by attenuating aortic degeneration. BIOMED PHARMACOTHER. 2023 Apr;160:114299 IHC; Rat . 36724640