

bs-0795R**[Primary Antibody]****Resistin Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000) Reactivity: Rat (predicted: Mouse) Predicted MW.: 12 kDa Subcellular Location: Secreted
Clonality: Polyclonal		
GeneID: 56729	SWISS: Q9HD89	
Target: Resistin		
Immunogen: KLH conjugated synthetic peptide derived from human Resistin: 21-114/114.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene belongs to the family defined by the mouse resistin-like genes. The characteristic feature of this family is the C-terminal stretch of 10 cys residues with identical spacing. The mouse homolog of this protein is secreted by adipocytes, and may be the hormone potentially linking obesity to type II diabetes. [provided by RefSeq]		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat breast); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Resistin) Polyclonal Antibody, Unconjugated (bs-0795R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=7.419]** Hirona Kugo. et al. Tricaprin can prevent the development of AAA by attenuating aortic degeneration. BIOMED PHARMACOTHER. 2023 Apr;160:114299 IHC ;Rat. 36724640