## bs-1180R

# [ Primary Antibody ]



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**RhoA Rabbit pAb** 

**GeneID: 387 SWISS:** P61586

Target: RhoA

**Immunogen:** KLH conjugated synthetic peptide derived from human RhoA:

101-193/193.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a member of the Rho family of small GTPases, which cycle between inactive GDP-bound and active GTP-bound states and function as molecular switches in signal transduction cascades. Rho proteins promote reorganization of the actin cytoskeleton and regulate cell shape, attachment, and motility. The protein encoded by this gene is prenylated at its C-terminus, and localizes to the cytoplasm and plasma membrane. It is thought to be important in cell locomotion. Overexpression of this gene is associated with tumor cell proliferation and metastasis. Multiple alternatively spliced variants, encoding the same protein, have been identified.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:200-800)


Reactivity: Human, Mouse, Rat

(predicted: Rabbit, Zebrafish, Chicken, Dog)


**Predicted** 21 kDa MW.:

**Subcellular Location:** Cell membrane ,Cytoplasm

### **VALIDATION IMAGES**



Paraformaldehyde-fixed, paraffin embedded (Mouse cerebellum); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RhoA) Polyclonal Antibody, Unconjugated (bs-1180R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RhoA) Polyclonal Antibody, Unconjugated (bs-1180R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

### SFI FCTFD CITATIONS -

- [IF=4.6] Jiaxu Zhou, et al. JTE-013 Alleviates Pulmonary Fibrosis by Affecting the RhoA/YAP Pathway and Mitochondrial Fusion/Fission. PHARMACEUTICALS-BASE. 2023 Oct;16(10):1444 IHC,IF,WB; Mouse. 37895915
- [IF=3.923] Pan T et al. MiR-128-1-5p regulates tight junction induced by selenium deficiency via targeting cell adhesion molecule 1 in broilers vein endothelial cells. J Cell Physiol. 2018 Nov;233(11):8802-8814. WB; broiler. 29904913
- [IF=3.448] Yu C et al. LncRNA AC009022. 1 enhances colorectal cancer cells proliferation, migration, and invasion by

kinase pathway." Chinese			al cell injury via Rho, .5214