bs-17804R

[Primary Antibody]

MRPL51 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500) IF (1:100-500)
GenelD: 51258	SWISS: Q4U2R6	ICC/IF (1:100-500)
Target: MRPL51		ELISA (1:5000-10000)
Immunogen: KLH conjugated synthetic peptide derived from human MRPL51: 21-128/128.		Reactivity: (predicted: Human, Mouse, Rat, Sheep, Cow, Horse)
Purification: affinity purified by	Protein A	
Concentration: 1mg/ml		Predicted
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: ^{12 kDa} Subcellular Location: ^{Cytoplasm}
nuclear genes and mitochondrion. M of a small 28S sub estimated 75% pr prokaryotic riboso difference betwee ribosomes is that species, the prote sequence, and so prevents easy reco encodes a putativ ribosomal protein	chondrial ribosomal proteins are encoded by I help in protein synthesis within the itochondrial ribosomes (mitoribosomes) consist unit and a large 39S subunit. They have an otein to rRNA composition compared to omes, where this ratio is reversed. Another in mammalian mitoribosomes and prokaryotic the latter contain a 5S rRNA. Among different ins comprising the mitoribosome differ greatly in metimes in biochemical properties, which ognition by sequence homology. This gene e 39S subunit protein and belongs to the L47P family. Pseudogenes corresponding to this gene mosomes 2p, 2q, 5p, and 10q. [provided by	