## bs-21842R

## [ Primary Antibody ]



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## - DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Cofilin Rabbit pAb

**GenelD:** 1072 **SWISS:** P23528

Target: Cofilin

Purification: affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** The protein encoded by this gene can polymerize and

depolymerize F-actin and G-actin in a pH-dependent manner. Increased phosphorylation of this protein by LIM kinase aids in Rho-induced reorganization of the actin cytoskeleton. Cofilin is a widely distributed intracellular actin-modulating protein that binds and depolymerizes filamentous F-actin and inhibits the polymerization of monomeric G-actin in a pH-dependent manner. It is involved in the translocation of actin-cofilin complex from

cytoplasm to nucleus.

**Applications: IHC-P** (1:100-500)

IHC-F (1:100-500) IF (1:100-500)

Reactivity: Mouse (predicted: Human,

Rat, Pig, Sheep, Cow,


Horse)

Predicted MW.: 20 kDa

Subcellular Cell membrane, Cytoplasm

Location: , Nucleus

## VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Cofilin) Polyclonal Antibody, Unconjugated (bs-21842R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.