


bs-8676R**[Primary Antibody]****IFI202 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**Target:** IFI202**Immunogen:** KLH conjugated synthetic peptide derived from mouse IFI-202: 121-220/445.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** IFI-202 is a primarily nuclear phosphoprotein which inhibits cell growth, in part by modulating transcriptional activity of NFkB, E2F, AP-1 and p53. Two related human proteins, MNDA (myeloid cell nuclear differentiation antigen) and IFI-16, have also been described. Expression of MNDA has been observed specifically in cells of the granulocyte-macrophage lineage. IFI-16 is constitutively expressed in various T and B cell lines and can be induced by IFN-g in HL60 cells. At least four of the Gene 200 cluster of IFN-inducible proteins, IFI-202, IFI-204, MNDA and IFI-16, are localized in the nucleus.**Applications:** **WB** (1:500-2000)
IHC-P (1:100-500)
IHC-F (1:100-500)
IF (1:100-500)**Reactivity:** Mouse, Rat**Predicted MW.:** 50 kDa**Subcellular Location:** Cytoplasm ,Nucleus**— VALIDATION IMAGES —**

Sample: Thymus (Mouse) Lysate at 40 ug
 Primary: Anti-IFI-202 (bs-8676R) at 1/300 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 50 kD
 Observed band size: 50 kD



Sample: Lymph (Mouse) Lysate at 40 ug Primary:
 Anti-IFI-202 (bs-8676R) at 1/300 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 50 kD
 Observed band size: 50 kD



Tissue/cell: rat brain tissue; 4%
 Paraformaldehyde-fixed and paraffin-
 embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-IFI-202 Polyclonal Antibody, Unconjugated(bs-8676R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining