

**bs-19640R****[ Primary Antibody ]****OLFML2B Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**


|   |   |  |
|---|---|--|
| <b>Host:</b> Rabbit<br><b>Clonality:</b> Polyclonal<br><b>GeneID:</b> 25903<br><b>Target:</b> OLFML2B<br><b>Immunogen:</b> KLH conjugated synthetic peptide derived from human OLFML2B: 451-550/750.<br><b>Purification:</b> affinity purified by Protein A<br><b>Concentration:</b> 1mg/ml<br><b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.<br>Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. | <b>Isotype:</b> IgG<br><b>SWISS:</b> Q68BL8 | <b>Applications:</b> <b>WB</b> (1:500-2000)<br><b>IHC-P</b> (1:100-500)<br><b>IHC-F</b> (1:100-500)<br><b>IF</b> (1:100-500)<br><b>Reactivity:</b> Human, Mouse<br>(predicted: Cow)<br><b>Predicted MW.:</b> 84 kDa<br><b>Subcellular Location:</b> Secreted |
|---|---|--|

**— VALIDATION IMAGES —**

Sample: Eye (Mouse) Lysate at 40 ug Epithelium (Mouse) Lysate at 40 ug Primary: Anti-OLFML2B(bs-19640R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 84kD Observed band size: 84kD



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (OLFML2B) Polyclonal Antibody, Unconjugated (bs-19640R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human cervical carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (OLFML2B) Polyclonal Antibody, Unconjugated (bs-19640R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

**— SELECTED CITATIONS —**

- **[IF=3.288]** Liu J et al. Bioinformatic exploration of OLFML2B overexpression in gastric cancer base on multiple analyzing tools.BMC Cancer. 2019 Mar 13;19(1):227. IHC ;Human. 30866865