

**bs-19647R****[ Primary Antibody ]****Opticin Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> ELISA (1:5000-10000)
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 26254	<b>SWISS:</b> Q9UBM4	
<b>Target:</b> Opticin		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human Opticin: 201-300/332.		
<b>Purification:</b> affinity purified by Protein A		<b>Reactivity:</b> (predicted: Human, Mouse, Rat, Rabbit, Cow, Dog, Cat, GuineaPig, Horse)
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		<b>Predicted MW.:</b> 37 kDa
<b>Background:</b> Opticin belongs to class III of the small leucine-rich repeat protein (SLRP) family. Members of this family are typically associated with the extracellular matrix. Opticin is present in significant quantities in the vitreous of the eye and also localizes to the cornea, iris, ciliary body, optic nerve, choroid, retina, and fetal liver. Opticin may noncovalently bind collagen fibrils and regulate fibril morphology, spacing, and organization. The opticin gene is mapped to a region of chromosome 1 that is associated with the inherited eye diseases age-related macular degeneration (AMD) and posterior column ataxia with retinosa pigmentosa (AXPC1). [provided by RefSeq, Jul 2008]		<b>Subcellular Location:</b> Secreted