bs-0310R-BF647

[Secondary Antibodies]

Rabbit Anti-Chicken IgG H&L, BF647 conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Rabbit Clonality: Polyclonal	lsotype: gG	Applications: IF (1:100-1000) Flow-Cyt (1:100-1000)
Target: Rabbit Anti-Chicken IgG H&L Purification: affinity purified by Protein A Concentration: 2.0 mg/ml		ICC/IF (1:100-1000) Excitation Spectrum: 647nm Emission spectrum: 666nm Reactivity: Chicken
glycerol.	vith 1% BSA, 0.03% Proclin300 and 50 year. Avoid repeated freeze/thaw cyo	
serum with normal le is important for our d molecules are produc adaptive immune res functions; to bind to recruit other cells and variability of the IgG g	gG), is one of the most abundant prot- vels between 8-17 mg/mL in adult blo efence against microorganisms and t ed by B lymphocytes as a part of our ponse. The IgG molecule has two sep the pathogen that elicited the respon- d molecules to destroy the antigen. Th pool is generated by somatic recombi ecificities in an individual at a given t be 1011 variants.	bod. IgG he arate se and to ne nation