

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## **GRM4** Rabbit pAb

Catalog Number: bs-1894R

Target Protein: GRM4
Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Mouse, Rat (predicted: Human, Rabbit, Pig)

Predicted MW: 102 kDa

Subcellular Cell membrane

Locations:

Entrez Gene: 2914

Swiss Prot: Q14833

**Source:** KLH conjugated synthetic peptide derived from human GRM4: 301-400/912.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: L glutamate is the major excitatory neurotransmitter in the central nervous system and


activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be

perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein coupled receptors, that have been divided into 3 groups on the basis of

sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic

AMP cascade but differ in their agonist selectivities.

## **VALIDATION IMAGES**



25 ug total protein per lane of various lysates (see on figure) probed with GRM4 polyclonal antibody, unconjugated (bs-1894R) at 1:1000 dilution and  $4^{\circ}$ C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.