

**bs-5357R****[ Primary Antibody ]****phospho-GABBR2 (Ser892) Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ELISA</b> (1:5000-10000)  <b>Reactivity:</b> Human (predicted: Mouse, Rat, Pig, Cow, Chicken, Dog)  <b>Predicted MW.:</b> 104 kDa  <b>Subcellular Location:</b> Cell membrane
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 9568	<b>SWISS:</b> O75899	
<b>Target:</b> GABBR2 (Ser892)		
<b>Immunogen:</b> KLH conjugated Synthesised phosphopeptide derived from human GABBR2 around the phosphorylation site of Ser892: RL(p-S)LQ.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> The multi-pass membrane protein encoded by this gene belongs to the G-protein coupled receptor 3 family and GABA-B receptor subfamily. The GABA-B receptors inhibit neuronal activity through G protein-coupled second-messenger systems, which regulate the release of neurotransmitters, and the activity of ion channels and adenylyl cyclase. This receptor subunit forms an active heterodimeric complex with GABA-B receptor subunit 1, neither of which is effective on its own. Allelic variants of this gene have been associated with nicotine dependence.[provided by RefSeq, Jan 2010].		

**— SELECTED CITATIONS —**

- **[IF=64.5]** Zhenyu Xiao. et al. 3D reconstruction of a gastrulating human embryo. CELL. 2024 Apr 23 IF ;Human. 38657603